

RESEARCH PAPER

Ethnopharmacological survey of medicinal plants in semi-arid rangeland in western Iran

Ehsan Zeidali ¹, Hosein Mardani Korrani ², Yaser Alizadeh ^{1*}, Fatemeh Kamari ³

¹ Department of Agronomy and Plant Breeding, Faculty of Agriculture, Ilam University, Ilam, Iran

² Department of Biological Production Science, Tokyo University of Agriculture and Technology, Tokyo, Japan


³ Department of Agronomy and Plant Breeding, Faculty of Agriculture, Zabol University, Zabol, Iran



Highlights

- In this study, a number of 61 native plant species was identified by medicinal use.
- The most of the collected native species by medicinal use were belonged to Asteraceae family (12 species).
- Fruit/ seed, Leaves and flower were recognized the main parts of plants those used as the traditional medicine in Chahmahal & Bakhtiari.

Graphical Abstract



Article Info

Receive Date: 21 November 2020

Revise Date: 21 January 2021

Accept Date: 18 February 2021

Available online:

Keywords:

Medicinal plants
Ethnopharmacological property
Chaharmahal and Bakhtiari province
Rangeland ecosystem

Abstract

Ethnopharmacology studies natural medicines derived from plants and other substances that have been traditionally used by groups of people to treat various human diseases. Various environmental conditions (from Tropical to Temperate condition) of Iran have made this country one of the major parts of plant diversity in the world. Plants have played an important role in Iranian people's life. The study is located in Chaharmahal and Bakhtiari province in western Iran (31°9' to 32°38' N and 49°30' to 51°26' E) with an area of 16,332 km² and a population of about 895,263. Endemic plants were collected during 2017-2019 from different locations in Chaharmahal and Bakhtiari province. Most of the studied regions were situated in orchards, highlands and neighboring farmlands. The plant raw materials were cleaned, dried and fixed at room temperature. In this study, a number of 61 native plant species with medicinal properties were collected during three years. Among the 70 people that were interviewed (mean age of 60 years old), 61 species of vascular plants were identified for treatment of various human ailments. Ethnopharmacology report is made consisting of species names, vernacular names, popular uses of the plants and their pharmacological properties. Identifying plant classification (taxa) was done in Chaharmahal and Bakhtiari Agricultural Research Center and the Iranian Research Institute of Plant Protection. The results showed that most of the collected species belonged to Asteraceae (12 species). Other families are included: Fabaceae (seven species), Brassicaceae (five species), Lamiaceae and Solanaceae (four species). This study shows the high herbal biodiversity of the medicinal plants in Iran as well as a fascinating potential for profitable studies on medicinal plant breeding, chemistry and pharmacology of the feature drugs.

© 2021 Published by CAS-Press.



doi:10.22034/CAJPSI.2021.01.03

E-ISSN: 2783-1310

* Corresponding author: yaseralizade5@gmail.com (Y. Alizadeh)

1. Introduction

Ethnopharmacology studies natural medicines derived from plants and other substances that have been traditionally used by groups of people to treat various human diseases. Ethnopharmacological studies of medicinal plants are useful in order to protect and find new uses of biological resources. Using medicinal plants to remedy human illness goes back to 2500 years ago which is transferred from elders to younger people orally, often from generation to generation. Recently, the subject of traditional medicine in Iran has received great attention in the literature, with regard to the use of medicinal plants (Rahman and Gulshana, 2014). The public tendency for the application of herbal and natural products for therapeutic purposes has enhanced in recent years. There are exciting possibilities for using plants as a source for the discovery of novel lead structures to be employed in drug industries, and also for the development of active plant extracts useful in treating a variety of ailments in humans and animals. The increasing global spread of drug's resistance to most of the available and affordable chemical drugs is a major apprehension that requires innovative strategies. Furthermore, it has been shown that more than 25% of chemical drugs are made of plants and 12% of them have microbial origins (Borris, 1996). Medicinal herbs have been introduced as the main source of the novel bio-molecule (Holland, 1994; Heinrich and Gibbons, 2001), which can theoretically be used in treating multiple lives-threatening illnesses such as malaria and cancer (Shoemaker et al., 2005; Cragg and Newman, 2005). Historical and cultural reasons are the other two factors responsible for the popularity of herbal medicines.

Various environmental conditions (from Tropical to Temperate condition) of Iran have made this country one of the major parts of plant diversity in the world. Plants have played an important role in Iranian people's life. Thousands of medicinal plants with potential pharmaceutical properties grow across the country. Moreover, germplasm and genetic resources are some of the more important natural wealth of any country. Documentation of the indigenous knowledge through ethnobotanical studies is important for the conservation and utilization of biological resources. Therefore, the establishment of the local names and indigenous uses of plants has significant potential societal benefits. There has been an extensive global attempt to identify medicinal components of plant material, but so far less than 10% of the total of 250000 plant species have been identified and explored for more than one biological function. Understanding genetic variation among populations is required as an effective practice for the conservation of endangered plants. Chaharmahal and Bakhtiari province is known as one of the most important regions of the medicinal plant diversity in Iran. A wealth of these plants is employed by the local people for the "treatment" of several ailments. For example, Persian shallot (*Allium hirtifolium*) bulbs to treat rheumatism, pain; Flixweed (*Descurainia Sophia*) to the cough and diarrhea; Chevill (*Ferulago subvelutina*) to cholesterol, uric acid and rheumatic; Keluss (*Kelussia odoratissima*) to the "fat blood," triglycerides and many other teas for many other disorders. Considering appropriate policies based on realistic consideration concerning the current position of natural ecosystems in some suitable practices and technical methods in all dimensions, including cultivation, managing, harvest and industrial and commercial exploitation of them either in natural habitats or in mechanized farms, needs a reliable approach to achieve a comprehensive view on the role and productivity of medicinal plants in developing countries like Iran.

The present study is carried out to assess and document the folk medicinal knowledge of the populations with respect to the wild area taxons. They are liable to become lost due, not just to the advances of civilization, but also to the aging of the local people and finding our common/rare uses of medicinal species, which could lead to the expansion of new plant-based medicines. Specifically, we aimed to answer the following questions of (i) what is the diversity of medicinal plants in the study area? (ii) What parts of the plant are used as medicine and what are their medicinal properties? (iii) What are the most important ailments in the area study and what plant species are used for their treatment?

2. Materials and Methods

2.1. Study area

The study is located in Chaharmahal and Bakhtiari province in western Iran (31°09' to 32°38' N and 49°30' to 51°26' E) with an area of 16,332 km² (Fig. 1) and a population of about 895,263. The annual precipitation is varied from less than 250 mm to more than 1000 mm. December and August are the coldest and the hottest months, respectively. Numerous varieties of wild plant species, due to different ecological conditions can therefore be found. Along with increased latitude, more dense vegetation is observed. Furthermore, the existence of some places with more than 3000 m altitudes in Koohrang is accompanied by the presence of some unique medicinal species such as *Kelussia odoratissima*, *Ferulago subvelutina* and *Echinophora platyloba*. Regions with different elevation levels in the study area have suitable environments for the growth and extension of plant species in the rangeland area. These lands are the valuable resources of plant species and medicinal herbs. Moreover, there are some kinds of rare plant species with medicinal properties and multiple uses, which grow as wild populations across the rangeland and farm margins such as Liquorice (*Glycyrrhiza glabra*), Flixweed (*Sisymbrium Sophia*) and Common Thyme (*Thymus vulgaris*).




Figure 1. Location of the study area on the map of Iran.

2.2. Plant collection

Endemic plants were collected during 2017-2019 from different locations in Chaharmahal and Bakhtiari province. Most of the studied regions were situated in orchards, highlands and neighboring farmlands. The plant raw materials were cleaned, dried and fixed at room temperature. The collected plants were identified by the Research Center of Agriculture and National Resource (RCANR) and Plant Pathology Research Institute (PPR) in Tehran.

A study in the context of interviewing was performed to investigate the traditional application of plants and to identify the parts of plants used by local people. Participants of the study composed of 65 locals well-known people of both genders with an average age of 65 years from the study area. Local people were asked to collect the name of the plants, to indicate against which illnesses the plants were used and to determine the methods of plant applications. Furthermore, only those species that could be directly indicated and collected by the local persons were studied. The vernacular names were collected with the cooperation of local people. For each species, the following information was collected: botanical taxon, family name, part used, common secondary metabolites, and references (Table 1).

3. Results and Discussion

According to the study results, it was observed that eight species had the highest concentration of the Alkaloids among which are *Hyoscyamus niger*, *Datura stramonium*, *Artemisia annua* and *Artemisia persica*. These species are important for pharmaceutical value and also for providing raw material. It was observed that 18 species were significant considering essential oils. Chamomile has a high level of necessary oils and flavonoids that have multiple applications in traditional medicine are *Echinophora platyloba*, *Ferulago subvelutina*, *Ziziphora clinopodioides*, *Artemisia annua* and *Tanacetum parthenium*. A number of 10 species were important regarding flavonoid compounds among which *Kelussia odoratissima* (also of essential oils) and *Hypericum perforatum* are the most important ones (Chalibian et al., 2006).

Considering these findings and the importance of providing raw material for pharmaceutical industries, conducting extensive future studies for investigation of morphological traits, genetic variation and breeding programs, especially for those plants exposed to extinction such as Khosharize (*Echinophora platyloba*), Mokhalase (*Tanacetum parthenium*) and Keluss (*Kelussia odoratissima*) which have not been investigated carefully seems to be necessary.

The parts used for treating ailments were included fruits/seeds (20.9%), leaves/petioles (17.7%), flowers (16.5%), stem/barks/wood (15.8%), underground parts (bulbs/rhizomes/tubers/roots) (13.9%), shoots (every parts above ground) (7.6%), and resins (3.8%). Leaves and barks were recognized as the main parts of plants used as the traditional maternal healthcare in Nigeria (Kankara et al., 2015). Leaves were recognized as the main part of plant species for preparing remedies (Kpodar et al., 2015). Furthermore, the whole plant was used in order to treat many ailments (3.8%). The most frequently used plant part was fruits/seeds (20.9%) followed by leaves/petioles (17.7%) (Fig. 2). Fruits/seeds and leaves/petioles are the most-used parts, possibly because they contain the higher amount of bioactive compounds than other parts (Rokaya et al., 2010). The harvest of the whole plant, fruits, seeds or roots affects negatively plant population growth, perhaps causing a strong decies of many medicinal plants in the natural ecosystems (Ghimire et al., 2008; Rokaya et al., 2010). In addition, the aerial parts such as leaves of the plants are maintained relatively more than other parts of the plants (Giday et al., 2003; Rokaya et al., 2010).

The results indicated that between 62 plant species with medicinal, the most frequently represented family were Asteraceae (19.36%) with 12 taxa, Fabaceae (11.29%), Brassicaceae (8.6%), Lamiaceae and Solanaceae (6.45%) (Fig. 2). Other representative families (11%) and Rosaceae (7%) are followed, respectively. Plant distribution across the province, exploited part of the plants, secondary metabolite, ethnopharmacology properties, the plant endemic and their scientific names are presented in Table 1. In this regard, investigation and comprehensive identification of the medicinal part of plants of the province needs more conceptual and financial investment from the scientific center (Table 1).

Based on the data reported by the Agricultural Research Center of Chaharmahal and Bakhtiari in 2010, the collection of plants obtained in the province is composed of 87 families, 468 genera and 1045 species, among which the most frequent genera belong to Asteraceae, Brassicaceae, and Poaceae with 69, 45 and 43 genera respectively. Among 1045 species identified in the province, 270 species possess pharmaceutical properties. The highest number of species is seen in Lamiaceae (35 species, 3.34%), Asteraceae (30 species, 2.87%), Apiaceae (20 species, 1.91%) and Fabaceae (19 species, 1.81%). Considering Raikaner's vegetative form, 46 Throphyt, 10 Cryptophyte, 119 Hemi-cryptophyt, 45 Phanerophyt, 45 Geophyt and 2 Parasite were identified. Regarding life form, 20 tree species 24 shrubs, 172 polyneal herbaceous, 7 biennial herbaceous and 3 annual occasionally biennial species have been detected.

The statistical population in this investigation covers 5.9% of the total flora of this province. As mentioned above, 270 (25.83%) species' pharmaceutical properties confirm the richness of the area in terms of medicinal plants. It was observed that fruit and seeds, leaves and petioles, flowers, stem, bark, wood and underground parts, with 20.9, 17.7, 16.5, 15.8, and 13.9% are the most utilized parts respectively (Fig. 2). Knowing the fact that utilizing underground parts such as root and tubers results in the eradication of the plants and that 22% of the

plants in this study are used for underground parts, extinction risk in these species is more serious and needs urgent attempt toward preventing their extinction.




Figure 2. Number of plant parts used for the treating ailment.

Table 1. Medicinal species traditionally used in Chaharmahal and Bakhtiari province in western Iran.

Scientific name (Family)	Commune secondary metabolite	Therapeutic properties	Parts used	Reference
<i>Acantholimon biois</i> (Caryophyllaceae)	Tannin, Amidon, Gallic acid	Carminative, Anti-hemintic, Astrigent, Healing	WP	Omidbaigi, 2009
<i>Achillea aleppica</i> (Asteraceae)	Azulene, Btonicine, Stachydrin, Mataicarin, Betaime, Luteolinn, Free Apigenin	Anti-inflammatory, regulates blood pressure, menstrual disorders, antipyretic, stimulating blood circulation	Fl, l	Omidbaigi, 2009; Zargari, 2003; Arihan et al., 2015
<i>Alcea angulate</i> (Malvaceae)	Mucilage, pectin, Amidon, acid valeric	anti-cough, respiratory tract emollient, antibacterial, Producing sputum	Fl, F	Omidbaigi, 2009; Zargari, 2003
<i>Alhagi persarum</i> (Fabaceae)	Fructose, Sucrose	Laxative, Expectorant, Febrifuge, Diuretic, Anti-Scabies	WP, Re	Shafizadeh, 2002
<i>Allium hirtifolium</i> (Alliaceae)	Sulfur compounds such as: allyl propyl sulfide	Diuretic, Diaphoretic, Expectorant, Anti-Parasit, Sedative, Aromatic, Flavor, Anti-rheumatism, Anti-core	B, L	Abbas et al., 2003 ; Bahmani et al., 2015; Taran et al., 2006; Jafarian et al., 2010
<i>Anagallis arvensis</i> (Primulaceae)	Saponin, Glicrilazide, Cyclamin	Icterus, Expectoran, Diuretic, Pruritis, Toothache, Anti-Pustules	WP	Shafizadeh, 2002
<i>Anthemis altissima</i> (Asteraceae)	Mucilage, chamazulene, luteolin-glycoside, apigenin, alfa bisabolol	Sedative, anti-inflammatory, analgesic menstruation, Memory Booster, booster digestive system	F	Acquaviva et al., 2012
<i>Arctium lappa</i> (Asteraceae)	Fatty acid, Oleic acid, Arctiol, Inulin, polyphenolic acid, Flavonoids	Diuretic, Diaphoretic, Stomachic, Cathartic, Febrifuge, Hypoglycemic	WP	Neves et al., 2007; Ziaie et al., 2006
<i>Artemisia annua</i> (Asteraceae)	Essential oils, Tannins, Inulins, Alkaloid, Mucilage	Stomachic, Diuretic, Diaphoretic, Astrigent, Vermifuge, Parasiticide	WP	Omidbaigi, 2009; Haynes and Krishna, 2004
<i>Artemisia persica</i> (Asteraceae)	Resin, Santonin, Artemisin, Terpens, Phllandrene, Thujone	Febrifuge, Stomachic, Disinfectant, Bechic, Vermifuge, Sadative	F	Omidbaigi, 2009

Table 1. Continued.

<i>Astragalus gossypinus</i> (Fabaceae)	Minerals, Saccharids, Gum	Laxative, Sedative, Hoarseness of voic, Use in preparation of drugs in emulsions and suspensions, Ophthalmia	Sy	Omidbaigi, 2009; Azadbakht, 2003
<i>Cannabis sativa</i> (Cannabaceae)	Cannabinol, Tetrahydrocannabinol, Cannin, Cannabinoid	Cathartic, Dysmenorrhea, Vomitting, Alopecia, Tonic,	F, L, Sh	Omidbaigi, 2009
<i>Capsella bursa pastoris</i> (Brassicaceae)	Tannin, Asetilkolin, Terpenoid, Resin, Myrosinase, Tyramine	Astringent, Diuretic, Hemostatic, Oligomenorrihea, Dysuria	WP	Shafiazadeh, 2002
<i>Carthamus oxyacantha</i> (Asteraceae)	Carthamin, Tartaric acid, Carthamul, Lignin, oil, Polysaccharids	Anti- Arteriosclerosis, Cathartic, Hypertensin, Abortive	F, Sd	Omidbaigi, 2009; Dini and Babakhanlou, 2002
<i>Centaurea luristanica</i> (Asteraceae)	Tannin, Cyanine, Pelargonien, glycosides	Febrifuge, Diuretic, Cathartic, Bechie, Expectorant, Eupeptic, Bronshit, Alopecia	F, L	Centaurea luristanica (Asteraceae)
<i>Cardaria draba</i> (Brassicaceae)	Essential oils, Ascorbic acid	Anti- scorbut, Breast Vulnerary, Hemorrhoid, Anti-core, Anti-dote	St, Br, L	Shafiazadeh, 2002
<i>Capparis spinosa</i> (Capparaceae)	Flavonoids, glycosides, glucocaparin	Stomachic, Expectorant, Diuretic, Emmenagogue, Toothache, Healing, Anti-dote, Parasiticide	F, R, Br, Fb	Karanayil et al., 2011; Shemshadi et al., 2015; Zeidali and Khorshidvand, 2015
<i>Cichorium intybus</i> (Asteraceae)	Tannin, Mucilage, Pectin, Inulin, Laktotropes, Sesquied, Potassium nitrate	Diuretic, Laxative, Diuphoretic, Anti-pruritis, Febrifuge, Pustules, Stomachic, Disinfectant, Choleretic, Mouth Wash	R, Fb, Sh, L	Haghi and Safaie, A, 2003; Zargari, 2003
<i>Citrullus colocynthis</i> (Cucurbitaceae)	Citrullol, Elaterin, Albuminoid, Coloyntetin, Alkaloid, Pectin	Backache, Headache, Cathartict, Emmenagogue, Epilepsy, Amnesia Toothache, Anti- rheumatism	F, Sd, R	Omidbaigi, 2009
<i>Convolvulus koeieanus</i> (Convolvulaceae)	Amidone, hydrocarbons	Choleretic, Hemostatic, Vulnerary, Healing	R, L, F, Sd	Shafiazadeh, 2002; Azadbakht, 2003
<i>Cynodon dactylon</i> (Poaceae)	Triticin, Inosit, Potassium	Diuretic, Diuphoretic, Anti- rheumatism, Anti- Pustules, Hypoglycemic	Rh	Singh et al., 2008
<i>Marrubium vulgare</i> (Lamiaceae)	Tannin, Mucilage, Lipids	Stomachic, Expectorant, Astringent, Febrifuge Emmenagogue, Anuria, Disinfectant, Choleretic	L, F, Sh	Zargari, 2003
<i>Melilotus officinalis</i> (Fabaceae)	Lysine, Cysteine, Phenylalanine, high level of vitamin(A, C, E, K)	Child Atony, stress, Stimulant, Anti-Migrena , Diuresis Menopause Hypotensor, Diuretic, Stomachic, Slimming	Sh, F, L, Sh	Melilotus officinalis (Fabaceae)
<i>Onopordum acanthium</i> (Asteraceae)	Oil, Inulin	Febrifuge, Diuretic, Astringent, Stomachic, Hemostatic, Anitifungi, Healing	Sd, R, Sh, F	Taleie et al., 2006
<i>Orobanche ramosa</i> (Orobanchaceae)	Sesquiterpene	Choleretic, Cornicid, Carminative, Anti-scorbutic, Laxative, Stomachic	St	Omidbaigi, 2009
<i>Ononis spinosa</i> (Fabaceae)	Glycosides, Ononin, Tannin, Citric acid	Diuretic, Anti-Scabies, Disinfectant	R, F, L	Zareazadeh et al., 2010
<i>Papaver rhoeas</i> (Papaveraceae)	Roemerine, Alkaloid, Papaveraldine, Codeine, Cryptopine, Tannin, Resin, Anthocyanin, Papaveric acid	Blepharritis, Sedative, Expectorant, Astringent, Hypnotic, Diuphoretic, Bechic, Hoarseness	F, Sd	Omidbaigi, 2009; Azadbakht, 2003

Table 1. Continued.

<i>Plantago psyllium</i> (<i>Plantaginaceae</i>)	Tannin, Tannic acid, Terpenes, Gallic acid, Octopine, Xylene, Mucilage, Fatty acid	Laxative, Expectorant, Febrifuge	Br, L, Sd	Singh and Chauhan, 2009
<i>Plantago lanceolata</i> (<i>Plantaginaceae</i>)	Mucilage, Tannin, Flavonoids, Saponozid, Ascorbic acid, Apigenin, Aucubin, Aucubin, Apigenin, AllAnti-on	Astringent, Anti-Anti-diarrhea, Hoarseness of voice, Anti- Pustules, Diuretic, Toothache	L, R, Sd	Zarian et al., 2004
<i>Peganum harmala</i> (<i>Zygophyllaceae</i>)	Alkaloids such as: Harmine, Harmaline, Harmalol	Hypnotic, Febrifuge, Diuretic, Parasiticide Emmenagogue, Anti- fungal Galactagogue, Toothache, Disinfectant	Sd, R, L	Zareazadeh et al., 2010; Diba et al., 2011; Omidbaigi, 2009
<i>Physalis divericata</i> (<i>Solanaceae</i>)	Ascorbic acid, Citric acid, Cryptoxantine, Physalin	Anti-helminthic, Diuretic, Emmenagogue, Sedative, Laxative, Anti-rheumatism, Carminative, Leterus	F	Dini and Babakhanlou, 2002
<i>Raphanus raphanistrum</i> (<i>Brassicaceae</i>)	Essential oils, Ascorbic acid, High levels of amino acid	Diuretic, Expectorant, Anti- influenza, Anti-Scurvy, Anti- rheumatism	Sy, L, R	Shafizadeh, 2002
<i>Reseda lutea</i> (<i>Resedaceae</i>)	Essential oil, Flavonoids, Glycosides	Diuretic, Diaphoretic, Cathartic, Stomachic	R, Sd, Sh, F	Diuretic, Diaphoretic, Cathartic, Stomachic
<i>Rheum ribes</i> (<i>Rubiaceae</i>)	Anthraquinone compounds, Reide, Malic acid, Funic acid, Sulfuric acid, Resin, Oxalic acid, Tannin, Antosianin	Stomachic, Emmenagogue, Astringent, Anti-lithiasis	Pt, R, Re	Zargari, 2003
<i>Sanguisorba minor</i> (<i>Rosaceae</i>)	Tannin, High levels of Minerals	Stomachic, Diuretic, Astringent, Anti-cold, Hemorrhoid, Oligomenorrhoea analgesic, astringent, anti-wrinkle, anti-viral,	Sanguisorba minor (<i>Rosaceae</i>)	Tannin, High levels of Minerals
<i>Solanum nigrum</i> (<i>Solanaceae</i>)	Solanine ,Saponin, Solasoidine Diosgenin, Tannin	Febrifuge, Hemostatic, Purgative, Toothache, Hemorrhoid	F, L	Azadbakht, 2003
<i>Silybum marianum</i> (<i>Asteraceae</i>)	Essential oils, high level of hydrocarbons	Anti-cirrosis	F, Br	Khalilian et al., 2003
<i>Salvia sclarea</i> (<i>Lamiaceae</i>)	Tannin, Phenic acid, Terpens, Acetone, Linalool, Salvion, Borneol	Anti- Hysteria, Febrifuge, Emmenagogue, Sedative, Blood suger, Toothache, Disinfectant, Analeptic, Hypotensor, Anti-core	L, Sh, F	Peana and Moretti, 2002
<i>Tanacetum persicum</i> (<i>Asteraceae</i>)	Essential oil, α -pinene, Terpenoid, Tannin, Flavonoid, Camphor	Anti-Hysteria, Emmenagogue, Astringent, Disinfectant, Febrifuge, Insecticide, Parasiticide, Migren	L, Sh, F	Mehrnia and Biranvand, 2007; Esmaeili et al., 2010
<i>Tragopogon caricifolius</i> (<i>Asteraceae</i>)	Inulin, protein, carbohydrate, minerals	Hypotensor, Hypoglycemic, Cholertic, Astringent, Hemostatic	F, L	Omidbaigi, 2009
<i>Tribulus terrestris</i> (<i>Zygophyllaceae</i>)	Cinnamic acid, Sterol, steroid, Amino acids, Alkaloid	Diuretic, Stomachic, Febrifuge, Astringent, Anti-scubies, Anti- lithiasis, hypotensor	L, Sh, F, R	Shafizadeh, 2002
<i>Astragalus hamosus</i> (<i>Fabaceae</i>)	Tannin, Coumarin, Flavin, Hexadecanoic acid	Expectorant, Carminative Astringent, Sedative, Headache, Vertigo, Amnesia	F, Sd, Sh	Azadbakht, 2003; Tawaha et al., 2007

Table 1. Continued.

<i>Datura stramonium</i> (<i>Solanaceae</i>)	Hyoscyamine, Atropin, Ascopolamin, Coumarin, Vitanol, Tannin	Headache, Febrifuge, Sedative, Aphasia, Simulative,	L, Sd, F	Yousaf et al., 2008; Aslani et al., 2007
<i>Urtica dioica</i> (<i>Urticaceae</i>)	Mucilage, Tannin, Asetilkolin, Secretin, Phytosterol, Gallic acid, Histamin, Glycerol, Ka+, Fe, Si	Diuretic, Diuphoretic, Astrigent, Pustules, Alopecia, Anti-goutal, Sedative, Hypoglycemic	L, Sd, R, Sy	Daher et al., 2006
<i>Vaccaria pyramidata</i> (<i>Caryophyllaceae</i>)	High level of protein, Saponin, Oil	Diuretic, Sedative, Anti-Parasit, Febrifuge, Anti-Scabies, Hemostatic, Dysmenorrhea	F, Sd, L	Omidbaigi, 2009; Zargari, 2003
<i>Verbascum lanceolatum</i> (<i>Scrophulariaceae</i>)	Saponin, Mucilage, Resin, Galactose, Arabinose	Diuretic, Diaphoretic , Anti- Hysteria, Expectorant, Emollient, Anti-asthma	F, L, R, Re	Omidbaigi, 2009; Zargari, 2003
<i>Veronica anagallis-aquatica</i> (<i>Scrophulariaceae</i>)	Tannin, Resin, Acobin, Wax	Stomachic, Tonic, Astrigent, Cicatrizing	F, L, St, Sh	Dini and Babakhanlou, 2002
<i>Viscum album</i> (<i>Loranthaceae</i>)	Viscotoxins Phoratoxin, Tannin, Terpenoid, Mucilage, Choline	Anti-oxidant, Diuretic, Hypotensor, Headache, Sedative	L, St, F	ÖnayUçar et al., 2006; Omidbaigi, 2009
<i>Ziziphora clinopodioides</i> (<i>Lamiaceae</i>)	Flavonoids, Carvacrol, Essential oil (Thymol), Tannin	Carminative, Hysteria, Ponic, Disinfectant, Diuphoretic, Expectorant, Melancholia, Migrene	L, Sh, St, F	Omidbaigi, 2009; Zargari, 2003

WP: Whole Plant, L: Leaves, St: Steam, Br: Bark, Re: Resin, F: Flower, R: Root, Sh: Shoot, Sy: Syrup

4. Conclusion

Promotional measures and strategies to prevent the increase of seed banks and vegetative organs in garden weed management for farmers should be explained. In general, research results show that weed density is high in Darreh Shahr gardens, so weed management methods should be improved to control weeds. Also, according to the results of this research, the structure of weeds in city gardens is different, and this can be useful in management planning for weed control.

References

- Acquaviva, R., Di Giacomo, C., Malaguarnera, M., Ragusa, S., Sorrenti, V., Santangelo, R., Mastrojeni, S., Barbagallo, I. Iauk, L., 2012. [Biological activities of extract of *Anthemis aetnensis* Schouw: In vitro evaluation. *J. Med. Plant. Res.*, 6\(10\), 1919-1925. <https://doi.org/10.5897/JMPR11.1567>](https://doi.org/10.5897/JMPR11.1567)
- Arihan, O., Özbek, H., Özkan, A.M.G., 2015. [Anti-inflammatory effects of *Sanguisorba minor* Scop. subsp. *muricata* \(Spach\) Briq. and *Cirsium libanoticum* DC. subsp. *lycaonicum* \(Boiss. & Heldr.\) Davis & Parris in rat. *East. J. Med.*, 20\(2\), 81-85. <https://doi.org/10.1016/j.ejme.2015.05.018>](https://doi.org/10.1016/j.ejme.2015.05.018)
- Aslani, M.R., Movasaghi, A., Janati, P.H., Karazma, M., 2007. [Experimental oleander \(*Nerium oleander*\) poisoning in goats: a clinical and pathological study. *Iran. J. Vet. Res.*, 8\(1\), 58-63. <https://doi.org/10.22099/ijvr.2007.2708>](https://doi.org/10.22099/ijvr.2007.2708)
- Bahmani, M., Saki, K., Ezatpour, B., Shahsavari, S., Eftekhari, Z., Jelodari, M., Rafieian Kopaei, M. Sepahvand, R., 2015. [Leishmaniosis phytotherapy: Review of plants used in Iranian traditional medicine on leishmaniasis. *Asian Pac. J. Trop. Biomed.*, 5\(9\), 695-701. <https://doi.org/10.1016/j.apjtb.2015.05.018>](https://doi.org/10.1016/j.apjtb.2015.05.018)
- Borris, R.P., 1996. [Natural products research: perspectives from a major pharmaceutical company. *J. Ethnopharmacol.*, 51\(1\), 29-38. \[https://doi.org/10.1016/0378-8741\\(95\\)01347-4\]\(https://doi.org/10.1016/0378-8741\(95\)01347-4\)](https://doi.org/10.1016/0378-8741(95)01347-4)

- Chalibian, F., Monfared, A., Larijani, K. Saldouzi, S., 2006. Comparison of the essential oils of *Chenopodium botrys* L., *Ferulago subvelutina* rech.F, *Rosa gallica* L. And antimicrobial activity of the oils against some microbes. *Iran. J. Med. Arom. Plant.*, **22**(2), 146-155. (In Persian) <https://doi.org/0.22092/ijmapr.2006.114989>
- Cragg, G.M., Newman, D.J., 2005. Plants as a source of anti-cancer agents. *J. Ethnopharmacol.*, **100**(1), 72-79. <https://doi.org/10.1016/j.jep.2005.05.011>
- Daher, C.F., Baroody, K.G., Baroody, G.M., 2006. Effect of *Urtica dioica* extract intake upon blood lipid profile in the rats. *Fitoterapia*, **77**(3), 183-188. <https://doi.org/10.1016/j.fitote.2006.01.010>
- Diba, K., Gerami Shoar, M., Shabatkhori, M., Khorshivand, Z., 2011. Anti fungal activity of alcoholic extract of *Peganum harmala* seeds. *J. Med. Plant. Res.*, **5**(23), 5550-5554. <https://doi.org/10.5897/JMPR.9000086>
- Dini, M., Babakhanlou, P., 2003. The checklist of useful plants. *Iran. J. Med. Arom. Plant*, **16**(1), 41-49. (In Persian)
- Esmaili, M.A., Sonboli, A., Noushabadi, M.A., 2010. Antioxidant and protective properties of six *Tanacetum* species against hydrogen peroxide-induced oxidative stress in K562 cell line: A comparative study. *Food Chem.*, **121**(1), 148-155. <https://doi.org/10.1016/j.foodchem.2009.12.022>
- Ghimire, S.K., Gimenez, O., Pradel, R., McKey, D., Aumeeruddy-Thomas, Y., 2008. Demographic variation and population viability in a threatened Himalayan medicinal and aromatic herb *Nardostachys grandiflora*: matrix modelling of harvesting effects in two contrasting habitats. *J. Appl. Ecol.*, **45**(1), 41-51. <https://doi.org/10.1111/j.1365-2664.2007.01375.x>
- Giday, M., Asfaw, Z., Elmqvist, T., Woldu, Z., 2003. An ethnobotanical study of medicinal plants used by the Zay people in Ethiopia. *J. Ethnopharmacol.*, **85**(1), 43-52. [https://doi.org/10.1016/S0378-8741\(02\)00359-8](https://doi.org/10.1016/S0378-8741(02)00359-8)
- Haynes, R.K., Krishna, S., 2004. Artemisinins: activities and actions. *Microbs. Infect.*, **6**(14), 1339-1346. <https://doi.org/10.1016/j.micinf.2004.09.002>
- Heinrich, M., Gibbons, S., 2001. Ethnopharmacology in drug discovery: an analysis of its role and potential contribution. *J. Pharm. Pharmacol.*, **53**(4), 425-432. <https://doi.org/10.1211/0022357011775712>
- Holland, B.K., 1994. Prospecting for drugs in ancient texts. *Nature*, **369**(6483), 702-702. <https://doi.org/10.1038/369702a0>
- Jafarian, A., Ghannadi, A., Elyasi, A., 2010. The effects of *Allium hirtifolium* Boiss. on cell-mediated immune response in mice. *Iran. J. Pharm. Res.*, **2**(1), 51-55. <https://doi.org/10.22037/ijpr.2010.36>
- Kankara, S.S., Ibrahim, M.H., Mustafa, M., Go, R., 2015. Ethnobotanical survey of medicinal plants used for traditional maternal healthcare in Katsina state, Nigeria. *South Afr. J. Bot.*, **97**, 165-175. <https://doi.org/10.1016/j.sajb.2015.01.007>
- Karanayil, R.S., Sinha, B.N., Rajasekaran, A., 2011. Protective Effects of *Capparis zeylanica* Linn. Leaf Extract on Gastric Lesions in Experimental Animals. *Avicenna J. Med. Biotechnol.*, **3**(1), 31-35.
- Kpodar, M.S., Lawson-Evi, P., Bakoma, B., Eklu-Gadegbeku, K., Agbonon, A., Aklikokou, K., Gbeassor, M., 2015. Ethnopharmacological survey of plants used in the treatment of diabetes mellitus in south of Togo (Maritime Region). *J. Herb. Med.*, **5**(3), 147-152. <https://doi.org/10.1016/j.hermed.2015.06.002>
- Neves, R.D.F., Moreno, S.R.F., Rebello, B.M., Caldas, L.Q.D.A., Fonseca, A.D.S.D., Bernardo-Filho, M., Medeiros, A.D.C., 2007. Effect of an *Arctium lappa* (burdock) extract on the labeling of blood constituents with technetium-99m and on the morphology of the red blood cells. *Braz. Arch. Biol. Technol.*, **50**(SPE) 167-174. <https://doi.org/10.1590/S1516-89132007000600021>
- ÖnayUçar, E., Karagöz, A., Arda, N., 2006. Antioxidant activity of *Viscum album* ssp. *album*. *Fitoterapia*, **77**(7-8), 556-560. <https://doi.org/10.1016/j.fitote.2006.08.001>
- Peana, A.T., Moretti, M.D., 2002. Pharmacological activities and applications of *Salvia sclarea* and *Salvia desoleana* essential oils. *Stud. Nat. Prod. Chem.*, **26**, 391-423. [https://doi.org/10.1016/S1572-5995\(02\)80012-6](https://doi.org/10.1016/S1572-5995(02)80012-6)
- Rahman, A.H.M.M., Gulshana, M.I.A., 2014. Taxonomy and medicinal uses on amaranthaceae family of Rajshahi, Bangladesh. *Appl. Ecol. Environ. Sci.*, **2**(2), 54-59. <https://doi.org/10.12691/aees-2-2-3>
- Rokaya, M.B., Münzbergová, Z., Timsina, B., 2010. Ethnobotanical study of medicinal plants from the Humla district of western Nepal. *J. Ethnopharmacol.*, **130**(3), 485-504. <https://doi.org/10.1016/j.jep.2010.05.036>

- Shemshadi, B., Ranjbar-Bahadory, S., Ahmadi, H., 2015. [Effect of Caparis spinosa root extract on promastigotes and amastigotes of Leishmania major](https://doi.org/10.22037/jps.v6i1.8053). *J. Paramed. Sci.*, **6**(1), 18-23. <https://doi.org/10.22037/jps.v6i1.8053>
- Shoemaker, M., Hamilton, B., Dairkee, S.H., Cohen, I., Campbell, M.J., 2005. [In vitro anticancer activity of twelve Chinese medicinal herbs](https://doi.org/10.1002/ptr.1702). *Phytother. Res.*, **19**(7), 649-651. <https://doi.org/10.1002/ptr.1702>
- Singh, B., Chauhan, N., 2009. [Modification of psyllium polysaccharides for use in oral insulin delivery](https://doi.org/10.1016/j.foodhyd.2008.06.004). *Food Hydrocoll.*, **23**(3), 928-935. <https://doi.org/10.1016/j.foodhyd.2008.06.004>
- Singh, S.K., Rai, P.K., Jaiswal, D., Rai, D., Sherma, B., Watal, G., 2008. [Protective effect of Cynodon dactylon against STZ induced hepatic injury in rats](https://doi.org/10.1016/j.jecp.2008.03.004). *J. Ecophysiol. Occup. Health*, **8**(3-4), 195-199.
- Taran, M., Rezaeian, M. and Izaddoost, M., 2006. [In vitro antitrichomonas activity of Allium hirtifloium \(Persian Shallot\) in comparison with metronidazole](https://doi.org/10.1016/j.jiph.2006.01.004). *Iran. J. Public Health*, **35**(1), 92-94.
- Tawaha, K., Alali, F.Q., Gharaibeh, M., Mohammad, M., El-Elimat, T., 2007. [Antioxidant activity and total phenolic content of selected Jordanian plant species](https://doi.org/10.1016/j.foodchem.2007.01.064). *Food Chem.*, **104**(4), 1372-1378. <https://doi.org/10.1016/j.foodchem.2007.01.064>
- Yousaf, Z., Masood, S., Shinwari, Z.K., Khan, M.A., Rabani, A., 2008. [Evaluation of taxonomic status of medicinal Species of the genus Hyoscyamos, Withania, Atropa and Datura based on poly acrylamide gel electrophoresis](https://doi.org/10.1016/j.pjbot.2008.06.004). *Pak. J. Bot.*, **40**(6), 2289-2297.
- Zareazadeh, A., Babakhanlou, P., Baghestani, N. and Shamszadeh, M., 2010. [Collection, identification and domestication of medicinal plant in Yazd province](https://doi.org/10.1016/j.jm.2010.03.004). *Iran. J. Med. Arom. Plant*, **14**, 73-135. (In Persian)



© 2020 by the authors. Submitted for possible open access publication under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

How to cite this paper:

Zeidali, E., Roein, Z., Fathi, A., 2020. [Ethnopharmacological survey of medicinal plants in semi-arid rangeland in western Iran](https://doi.org/10.22037/jps.v1i1.4655). *Cent. Asian J. Plant Sci. Innov.*, **1**(1), 46-55.